

Harel EH, Brown WD Attention deficit hyperactivity disorder in elementary school children in Rhode Island: associated psychosocial factors and medications used.

Clin Pediatr (Phila). 2003 Jul-Aug;42(6):497-503.

.Charach A¹, Cao H, Schachar R, To T Correlates of methylphenidate use in Canadian children: a cross-sectional study.

Can J Psychiatry. 2006 Jan;51(1):17-26

Strohschein LA. Prevalence of methylphenidate use among Canadian children following parental divorce. *CMAJ: Canadian Medical Association Journal.* 2007;176(12):1711-1714. doi:10.1503/cmaj.061458.

ELLIS B, NIGG J. Parenting Practices and Attention-Deficit/Hyperactivity Disorder: Partial Specificity of Effects. *Journal of the American Academy of Child and Adolescent Psychiatry.* 2009;48(2):146-154. doi:10.1097/CHI.0b013e31819176d0.

Sawyer, M. G., Rey, J. M., Graetz, B. W., Clark, J. J., & Baghurst, P. A. (2002). Use of medication by young people with attention-deficit/hyperactivity disorder. *Medical journal of Australia, 177*(1), 21-25.

Joseph, J. (2000). Not in their genes: A critical view of the genetics of attention-deficit hyperactivity disorder. *Developmental Review, 20*(4), 539-567.

Baumrind, D. (2005), Patterns of parental authority and adolescent autonomy. *New Directions for Child and Adolescent Development, 2005:* 61–69. doi:10.1002/cd.128